



Contents lists available at ScienceDirect

# Journal of Mathematical Analysis and Applications

[www.elsevier.com/locate/jmaa](http://www.elsevier.com/locate/jmaa)


## A Voronovskaya Type Theorem for Poisson–Cauchy Type singular operators

George A. Anastassiou\*, Razvan A. Mezei

Department of Mathematical Sciences, The University of Memphis, Memphis, TN 38152, USA

### ARTICLE INFO

#### Article history:

Received 23 June 2009

Available online 18 January 2010

Submitted by R. Curto

#### Keywords:

Smooth Poisson–Cauchy Type singular

integral

Voronovskaya Type Theorem

Rate of convergence

### ABSTRACT

In this article we continue with the study of approximation properties of smooth Poisson–Cauchy Type singular integral operators over the real line. We produce Voronovskaya Type results and give some quantitative results regarding the rate of convergence of these singular integral operators.

© 2010 Elsevier Inc. All rights reserved.

### 1. Introduction

We are motivated by the approximation properties of *Poisson–Cauchy Type singular integral* of a function  $f$  defined by the following

$$M_{\xi}(f; x) := \frac{\Gamma(\beta)\alpha\xi^{2\alpha\beta-1}}{\Gamma(\frac{1}{2\alpha})\Gamma(\beta - \frac{1}{2\alpha})} \int_{-\infty}^{\infty} f(x+y) \frac{1}{(y^{2\alpha} + \xi^{2\alpha})^{\beta}} dy, \quad (1)$$

for all  $x \in \mathbb{R}$ ,  $\xi > 0$ ,  $\alpha \in \mathbb{N}$ ,  $\beta > \frac{1}{2\alpha}$ , see [11], and see also [5, Chps. 16, 17], [4, Chp. 21], [6]. Other motivations come from [1–3, 7–10, 12, 14].

Next we define the *smooth Poisson–Cauchy Type singular integral operators*  $M_{r,n,\xi}(f; x)$  of which basic approximation properties were studied in [11].

For  $r \in \mathbb{N}$  and  $n \in \mathbb{Z}_+$  we set

$$\alpha_j = \begin{cases} (-1)^{r-j} \binom{r}{j} j^{-n}, & j = 1, \dots, r, \\ 1 - \sum_{j=1}^r (-1)^{r-j} \binom{r}{j} j^{-n}, & j = 0, \end{cases} \quad (2)$$

that is  $\sum_{j=0}^r \alpha_j = 1$ .

Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be such that  $f^{(n)}$  exists and it is bounded and Lebesgue measurable. We define for  $x \in \mathbb{R}$ ,  $\xi > 0$  and  $\alpha \in \mathbb{N}$ ,  $\beta > \frac{1}{2\alpha}$  the Lebesgue singular integrals

$$M_{r,n,\xi}(f; x) = \frac{\Gamma(\beta)\alpha\xi^{2\alpha\beta-1}}{\Gamma(\frac{1}{2\alpha})\Gamma(\beta - \frac{1}{2\alpha})} \int_{-\infty}^{\infty} \sum_{j=0}^r \alpha_j f(x+jt) \frac{1}{(t^{2\alpha} + \xi^{2\alpha})^{\beta}} dt. \quad (3)$$

\* Corresponding author.

E-mail addresses: [ganastss@memphis.edu](mailto:ganastss@memphis.edu) (G.A. Anastassiou), [rmezei@memphis.edu](mailto:rmezei@memphis.edu) (R.A. Mezei).

**Note 1.** The operators  $M_{r,n,\xi}$  are not in general positive, see [11].

**Note 2.** In particular we have  $M_{1,n,\xi} = M_\xi$ .

In Section 2 we give an elementary property of integrals related to (3). Then in Section 3, we will prove Voronovskaya type asymptotic results, see also [13].

## 2. An auxiliary result

From (3) we also obtain

$$M_{r,n,\xi}(f; x) = \frac{\Gamma(\beta)\alpha\xi^{2\alpha\beta-1}}{\Gamma(\frac{1}{2\alpha})\Gamma(\beta - \frac{1}{2\alpha})} \sum_{j=0}^r \alpha_j \int_{-\infty}^{\infty} f(x+jt) \frac{1}{(t^{2\alpha} + \xi^{2\alpha})^\beta} dt. \quad (4)$$

By means of elementary calculations, we obtain

**Lemma 3.** For every  $\alpha \in \mathbb{N}$ ,  $n \in \mathbb{N}_0$ ,  $\beta > \frac{n+1}{2\alpha}$ , and  $\xi > 0$ , we have

$$I_n := \int_0^\infty \frac{t^n}{(t^{2\alpha} + \xi^{2\alpha})^\beta} dt = \frac{1}{2\alpha\xi^{2\alpha\beta-n-1}} \frac{\Gamma(\frac{n+1}{2\alpha})\Gamma(\beta - \frac{n+1}{2\alpha})}{\Gamma(\beta)}. \quad (5)$$

**Proof.** See [15, p. 397, formula 595].  $\square$

## 3. Results

We present our main result.

**Theorem 4.** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be such that  $f^{(n)}$  exists,  $n \in \mathbb{N}$ , and is bounded and Lebesgue measurable on  $\mathbb{R}$ , and let  $\xi \rightarrow 0+$ ,  $0 < \gamma \leq 1$ ,  $\alpha \in \mathbb{N}$ ,  $\beta > \frac{n+1}{2\alpha}$ . Then

$$M_{r,n,\xi}(f; x) - f(x) = \sum_{m=1}^{\lfloor \frac{n-1}{2} \rfloor} f^{(2m)}(x) \left( \sum_{j=1}^r \alpha_j j^{2m} \right) \frac{\Gamma(\frac{2m+1}{2\alpha})\Gamma(\beta - \frac{2m+1}{2\alpha})}{\Gamma(\frac{1}{2\alpha})\Gamma(\beta - \frac{1}{2\alpha})} \frac{\xi^{2m}}{(2m)!} + o(\xi^{n-\gamma}). \quad (6)$$

**Proof.** We notice by  $\frac{\Gamma(\beta)\alpha\xi^{2\alpha\beta-1}}{\Gamma(\frac{1}{2\alpha})\Gamma(\beta - \frac{1}{2\alpha})} \int_{-\infty}^{\infty} \frac{1}{(t^{2\alpha} + \xi^{2\alpha})^\beta} dt = 1$ , that  $M_{r,n,\xi}(c, x) = c$ , for any  $c$  constant, and therefore we have

$$M_{r,n,\xi}(f; x) - f(x) = \frac{\Gamma(\beta)\alpha\xi^{2\alpha\beta-1}}{\Gamma(\frac{1}{2\alpha})\Gamma(\beta - \frac{1}{2\alpha})} \left( \sum_{j=0}^r \alpha_j \int_{-\infty}^{\infty} (f(x+jt) - f(x)) \frac{1}{(t^{2\alpha} + \xi^{2\alpha})^\beta} dt \right). \quad (7)$$

Using Taylor's formula for  $f$ , we have

$$f(x+jt) = \sum_{k=0}^{n-1} \frac{f^{(k)}(x)}{k!} (jt)^k + \frac{f^{(n)}(\theta)}{n!} (jt)^n, \quad (8)$$

with  $\theta = \theta(j)$  between  $x$  and  $x+jt$ .

We obtain

$$\begin{aligned} M_{r,n,\xi}(f; x) - f(x) &\stackrel{(7)}{=} \frac{\Gamma(\beta)\alpha\xi^{2\alpha\beta-1}}{\Gamma(\frac{1}{2\alpha})\Gamma(\beta - \frac{1}{2\alpha})} \\ &\quad \times \left( \sum_{j=1}^r \alpha_j \int_{-\infty}^{\infty} \left( \left[ \sum_{k=0}^{n-1} \frac{f^{(k)}(x)}{k!} (jt)^k + \frac{f^{(n)}(\theta)}{n!} (jt)^n \right] - f(x) \right) \frac{1}{(t^{2\alpha} + \xi^{2\alpha})^\beta} dt \right) \\ &= \frac{\Gamma(\beta)\alpha\xi^{2\alpha\beta-1}}{\Gamma(\frac{1}{2\alpha})\Gamma(\beta - \frac{1}{2\alpha})} \left( \sum_{j=1}^r \alpha_j \int_{-\infty}^{\infty} \left[ \sum_{k=1}^{n-1} \frac{f^{(k)}(x)}{k!} j^k t^k + \frac{f^{(n)}(\theta)}{n!} j^n t^n \right] \frac{1}{(t^{2\alpha} + \xi^{2\alpha})^\beta} dt \right) \end{aligned}$$

$$\begin{aligned}
&= \frac{\Gamma(\beta)\alpha\xi^{2\alpha\beta-1}}{\Gamma(\frac{1}{2\alpha})\Gamma(\beta-\frac{1}{2\alpha})} \left( \sum_{j=1}^r \alpha_j \left[ \sum_{k=1}^{n-1} \frac{f^{(k)}(x)}{k!} j^k \left( \int_{-\infty}^{\infty} \frac{t^k}{(t^{2\alpha} + \xi^{2\alpha})^\beta} dt \right) \right. \right. \\
&\quad \left. \left. + \frac{j^n}{n!} \int_{-\infty}^{\infty} f^{(n)}(\theta) t^n \frac{1}{(t^{2\alpha} + \xi^{2\alpha})^\beta} dt \right] \right) \\
&\stackrel{(5)}{=} \frac{\Gamma(\beta)\alpha\xi^{2\alpha\beta-1}}{\Gamma(\frac{1}{2\alpha})\Gamma(\beta-\frac{1}{2\alpha})} \left( \sum_{j=1}^r \alpha_j \left[ \sum_{m=1}^{\lfloor \frac{n-1}{2} \rfloor} \frac{f^{(2m)}(x)}{(2m)!} j^{2m} \frac{1}{\alpha\xi^{2\alpha\beta-2m-1}} \frac{\Gamma(\frac{2m+1}{2\alpha})\Gamma(\beta-\frac{2m+1}{2\alpha})}{\Gamma(\beta)} \right. \right. \\
&\quad \left. \left. + \frac{j^n}{n!} \int_{-\infty}^{\infty} f^{(n)}(\theta) t^n \frac{1}{(t^{2\alpha} + \xi^{2\alpha})^\beta} dt \right] \right) \\
&= \sum_{j=1}^r \alpha_j \left[ \left( \sum_{m=1}^{\lfloor \frac{n-1}{2} \rfloor} f^{(2m)}(x) \frac{\Gamma(\frac{2m+1}{2\alpha})\Gamma(\beta-\frac{2m+1}{2\alpha})}{\Gamma(\frac{1}{2\alpha})\Gamma(\beta-\frac{1}{2\alpha})} \frac{j^{2m}\xi^{2m}}{(2m)!} \right) \right. \\
&\quad \left. + \frac{\Gamma(\beta)\alpha\xi^{2\alpha\beta-1}}{\Gamma(\frac{1}{2\alpha})\Gamma(\beta-\frac{1}{2\alpha})} \frac{j^n}{n!} \int_{-\infty}^{\infty} f^{(n)}(\theta) t^n \frac{1}{(t^{2\alpha} + \xi^{2\alpha})^\beta} dt \right].
\end{aligned}$$

Therefore we have obtained

$$\begin{aligned}
M_{r,n,\xi}(f; x) - f(x) - \sum_{m=1}^{\lfloor \frac{n-1}{2} \rfloor} f^{(2m)}(x) \left( \sum_{j=1}^r \alpha_j j^{2m} \right) \frac{\Gamma(\frac{2m+1}{2\alpha})\Gamma(\beta-\frac{2m+1}{2\alpha})}{\Gamma(\frac{1}{2\alpha})\Gamma(\beta-\frac{1}{2\alpha})} \frac{\xi^{2m}}{(2m)!} \\
= \frac{\Gamma(\beta)\alpha\xi^{2\alpha\beta-1}}{\Gamma(\frac{1}{2\alpha})\Gamma(\beta-\frac{1}{2\alpha})} \sum_{j=1}^r \alpha_j \frac{j^n}{n!} \int_{-\infty}^{\infty} f^{(n)}(\theta) t^n \frac{1}{(t^{2\alpha} + \xi^{2\alpha})^\beta} dt.
\end{aligned} \tag{9}$$

Hence

$$\begin{aligned}
\Delta_\xi &:= \frac{1}{\xi^n} \left[ (M_{r,n,\xi}(f; x) - f(x)) - \sum_{m=1}^{\lfloor \frac{n-1}{2} \rfloor} f^{(2m)}(x) \left( \sum_{j=1}^r \alpha_j j^{2m} \right) \frac{\Gamma(\frac{2m+1}{2\alpha})\Gamma(\beta-\frac{2m+1}{2\alpha})}{\Gamma(\frac{1}{2\alpha})\Gamma(\beta-\frac{1}{2\alpha})} \frac{\xi^{2m}}{(2m)!} \right] \\
&= \frac{\Gamma(\beta)\alpha\xi^{2\alpha\beta-1-n}}{\Gamma(\frac{1}{2\alpha})\Gamma(\beta-\frac{1}{2\alpha})} \sum_{j=1}^r \alpha_j \frac{j^n}{n!} \int_{-\infty}^{\infty} f^{(n)}(\theta) t^n \frac{1}{(t^{2\alpha} + \xi^{2\alpha})^\beta} dt \\
&= \frac{\Gamma(\beta)\alpha\xi^{2\alpha\beta-1-n}}{n!\Gamma(\frac{1}{2\alpha})\Gamma(\beta-\frac{1}{2\alpha})} \int_{-\infty}^{\infty} \left( \sum_{j=1}^r \alpha_j j^n f^{(n)}(\theta) \right) \frac{t^n}{(t^{2\alpha} + \xi^{2\alpha})^\beta} dt \\
&= \frac{\Gamma(\beta)\alpha\xi^{2\alpha\beta-1-n}}{n!\Gamma(\frac{1}{2\alpha})\Gamma(\beta-\frac{1}{2\alpha})} \left[ \int_{-\infty}^{\infty} \left( \sum_{j=1}^r (-1)^{r-j} \binom{r}{j} f^{(n)}(\theta) \right) \frac{t^n}{(t^{2\alpha} + \xi^{2\alpha})^\beta} dt \right].
\end{aligned} \tag{10}$$

Call

$$\Phi_n(x, t) := \sum_{j=1}^r (-1)^{r-j} \binom{r}{j} f^{(n)}(\theta). \tag{11}$$

Thus

$$\Delta_\xi = \frac{\Gamma(\beta)\alpha\xi^{2\alpha\beta-1-n}}{n!\Gamma(\frac{1}{2\alpha})\Gamma(\beta-\frac{1}{2\alpha})} \left[ \int_{-\infty}^{\infty} \Phi_n(x, t) \frac{t^n}{(t^{2\alpha} + \xi^{2\alpha})^\beta} dt \right]. \tag{12}$$

Since we assumed that  $f^{(n)}$  exists and it is bounded we obtain

$$\|f^{(n)}\|_\infty < M, \quad \text{for some } M \geq 0.$$

Therefore

$$\begin{aligned} |\Phi_n(x, t)| &\leq \left( \sum_{j=1}^r \binom{r}{j} \right) M \\ &= (2^r - 1)M. \end{aligned} \quad (13)$$

Hence

$$|\Delta_\xi| \leq \frac{(2^r - 1)M \Gamma(\frac{n+1}{2\alpha}) \Gamma(\beta - \frac{n+1}{2\alpha})}{n! \Gamma(\frac{1}{2\alpha}) \Gamma(\beta - \frac{1}{2\alpha})} =: \lambda. \quad (14)$$

Consequently we get ( $0 < \gamma \leq 1$ )

$$\frac{1}{\xi^{n-\gamma}} \left| \left( M_{r,n,\xi}(f; x) - f(x) \right) - \sum_{m=1}^{\lfloor \frac{n-1}{2} \rfloor} f^{(2m)}(x) \left( \sum_{j=1}^r \alpha_j j^{2m} \right) \frac{\Gamma(\frac{2m+1}{2\alpha}) \Gamma(\beta - \frac{2m+1}{2\alpha})}{\Gamma(\frac{1}{2\alpha}) \Gamma(\beta - \frac{1}{2\alpha})} \frac{\xi^{2m}}{(2m)!} \right| \leq \lambda \xi^\gamma \rightarrow 0, \quad (15)$$

as  $\xi \rightarrow 0+$ .

Notice that  $n - 1 - 2m \geq 0$ .

From the last we conclude the claim of the theorem.  $\square$

**Corollary 5** ( $n = 1$  case). Let  $f$  such that  $f'$  exists and is bounded and Lebesgue measurable on  $\mathbb{R}$ . Let  $\xi \rightarrow 0+$ ,  $0 < \gamma \leq 1$ ,  $\alpha \in \mathbb{N}$ ,  $\beta > \frac{1}{\alpha}$ . Then

$$M_{r,1,\xi}(f; x) - f(x) = o(\xi^{1-\gamma}). \quad (16)$$

**Proof.** In Theorem 4 we place  $n = 1$ .  $\square$

**Corollary 6** ( $n = 2$  case). Let  $f$  such that  $f''$  exists and is bounded and Lebesgue measurable on  $\mathbb{R}$ . Let  $\xi \rightarrow 0+$ ,  $0 < \gamma \leq 1$ ,  $\alpha \in \mathbb{N}$ ,  $\beta > \frac{3}{2\alpha}$ . Then

$$M_{r,2,\xi}(f; x) - f(x) = o(\xi^{2-\gamma}). \quad (17)$$

**Proof.** In Theorem 4 we place  $n = 2$ .  $\square$

**Corollary 7** ( $n = 3$  case). Let  $f$  such that  $f^{(3)}$  exists and is bounded and Lebesgue measurable on  $\mathbb{R}$ . Let  $\xi \rightarrow 0+$ ,  $0 < \gamma \leq 1$ ,  $\alpha \in \mathbb{N}$ ,  $\beta > \frac{\alpha}{2}$ . Then

$$M_{r,3,\xi}(f; x) - f(x) = \xi^2 f''(x) \left( \sum_{j=1}^r \alpha_j j^2 \right) \frac{\Gamma(\frac{3}{2\alpha}) \Gamma(\beta - \frac{3}{2\alpha})}{2\Gamma(\frac{1}{2\alpha}) \Gamma(\beta - \frac{1}{2\alpha})} + o(\xi^{3-\gamma}). \quad (18)$$

**Proof.** In Theorem 4 we place  $n = 3$ .  $\square$

**Corollary 8** ( $n = 4$  case). Let  $f$  such that  $f^{(4)}$  exists and is bounded and Lebesgue measurable on  $\mathbb{R}$ . Let  $\xi \rightarrow 0+$ ,  $0 < \gamma \leq 1$ ,  $\alpha \in \mathbb{N}$ ,  $\beta > \frac{5}{2\alpha}$ . Then

$$M_{r,4,\xi}(f; x) - f(x) = \xi^2 f''(x) \left( \sum_{j=1}^r \alpha_j j^2 \right) \frac{\Gamma(\frac{3}{2\alpha}) \Gamma(\beta - \frac{3}{2\alpha})}{2\Gamma(\frac{1}{2\alpha}) \Gamma(\beta - \frac{1}{2\alpha})} + o(\xi^{4-\gamma}). \quad (19)$$

**Proof.** In Theorem 4 we place  $n = 4$ .  $\square$

## References

- [1] G.A. Anastassiou, Basic convergence with rates of smooth Picard singular operators, J. Comput. Anal. Appl. 8 (4) (2006) 313–334.
- [2] G.A. Anastassiou,  $L_p$  convergence with rates of smooth Picard singular operators, in: Differential & Difference Equations and Applications, Hindawi Publ. Corp., New York, 2006, pp. 31–45.
- [3] G.A. Anastassiou, Global smoothness and uniform convergence of smooth Picard singular operators, Comput. Math. Appl. 50 (10–12) (2005) 1755–1766.
- [4] G.A. Anastassiou, Quantitative Approximations, Chapman & Hall/CRC, Boca Raton, New York, 2001.
- [5] G.A. Anastassiou, S.G. Gal, Approximation Theory, Birkhäuser, Boston, Berlin, 2000.
- [6] G.A. Anastassiou, S. Gal, Convergence of generalized singular integrals to the unit, multivariate case, in: Appl. Math. Rev., vol. 1, World Sci. Publ. Co., Singapore, 2000, pp. 1–8.

- [7] G.A. Anastassiou, R.A. Mezei,  $L_p$  convergence with rates of smooth Gauss–Weierstrass singular operators, *Nonlinear Stud.*, in press.
- [8] G.A. Anastassiou, R.A. Mezei, Uniform convergence with rates of smooth Gauss–Weierstrass singular integral operators, *Appl. Anal.* 88 (7) (2009) 1015–1037.
- [9] G.A. Anastassiou, R.A. Mezei, Global smoothness and uniform convergence of smooth Gauss–Weierstrass singular operators, *Math. Comput. Modelling* (2009), doi:10.1016/j.mcm.2009.04.001.
- [10] G.A. Anastassiou, R.A. Mezei, Voronovskaya type theorems for smooth Picard and Gauss–Weierstrass singular operators, submitted for publication.
- [11] G.A. Anastassiou, R.A. Mezei, Global smoothness and uniform convergence of smooth Poisson–Cauchy type singular operators, *Appl. Math. Comput.* 215 (2009) 1718–1731.
- [12] R.A. DeVore, G.G. Lorentz, *Constructive Approximation*, vol. 303, Springer-Verlag, Berlin, New York, 1993.
- [13] L. Rempulska, Z. Walczak, On modified Picard and Gauss–Weierstrass singular integrals, *Ukrain. Mat. Zh.* 57 (11) (2005) 1577–1584; translation in: *Ukrainian Math. J.* 57 (11) (2005) 1844–1852.
- [14] L. Schumaker, *Spline Functions. Basic Theory*, J. Wiley & Sons, New York, 1981.
- [15] D. Zwillinger, *CRC Standard Mathematical Tables and Formulae*, 30th edition, Chapman & Hall/CRC, Boca Raton, 1995.